

Supplementary Table 1. NAFLD index for predicting MACCE using the Cox-regression analysis

Variable	Hazard ratios (95% confidence interval)			
	Unadjusted	Model 1	Model 2	Model 3
Hepatic steatosis index				
Cut-off value for detecting NAFLD ^a				
< 30	Reference	Reference	Reference	Reference
30–36	1.049 (0.908–1.213)	1.251 (1.079–1.450)	1.149 (0.956–1.382)	1.077 (0.895–1.296)
> 36	1.222 (1.039–1.437)	1.568 (1.326–1.854)	1.307 (0.998–1.713)	1.143 (0.871–1.502)
Fatty liver index				
Cut-off value for detecting NAFLD ^a				
< 30	Reference	Reference	Reference	Reference
30–<60	1.519 (1.329–1.737)	1.382 (1.209–1.581)	1.428 (1.244–1.639)	1.304 (1.133–1.501)
≥ 60	1.809 (1.556–2.103)	1.748 (1.501–2.036)	1.796 (1.533–2.104)	1.569 (1.334–1.846)
NAFLD liver fat score				
Cut-off value for detecting NAFLD ^a				
> -0.640	1.726 (1.508–1.976)	1.558 (1.360–1.784)	1.376 (1.162–1.628)	1.270 (1.073–1.504)
Fibrosis-4 index				
Cut-off value for detecting fibrosis in NAFLD ^a				
< 1.3	Reference	Reference	Reference	Reference
1.3–2.67	1.737 (1.533–1.970)	0.971 (0.847–1.113)	1.030 (0.894–1.185)	1.058 (0.919–1.218)
> 2.67	2.198 (1.573–3.073)	1.115 (0.792–1.569)	1.076 (0.748–1.547)	1.076 (0.748–1.547)

NAFLD, nonalcoholic fatty liver disease; MACCE, major adverse cardiovascular and cerebrovascular events.

^aCut-off value for detecting hepatic steatosis or liver fibrosis as suggested in the NAFLD guideline [34,39–41].