

Supplementary Table 1. Comparison of each component of the current perception threshold values between sleep quality groups

| Variable | Painless DPN (n=76) | No DPN (n=70) | P value |
|-----------------|------------------------|------------------|---------|
| 2,000 Hz | | | |
| Hand-left | 320.8±90.9 | 259.1±55.0 | <0.001 |
| Hand-right | 284.9±103.4 | 269.4±51.3 | 0.249 |
| Foot-left | 347.1±187.6 | 304.6±76.9 | 0.072 |
| Foot-right | 358.8±178.8 | 294.1±79.9 | 0.005 |
| 250 Hz | | | |
| Hand-left | 146.5±108.1 | 115.8±32.5 | 0.024 |
| Hand-right | 123.3±126.3 | 114.3±29.2 | 0.547 |
| Foot-left | 151.6±152.2 | 124.5±37.6 | 0.137 |
| Foot-right | 158.9±159.2 | 123.9±34.9 | 0.065 |
| 5 Hz | | | |
| Hand-left | 84.6±56.2 | 71.9±23.4 | 0.072 |
| Hand-right | 77.9±120.1 | 64.0±25.8 | 0.327 |
| Foot-left | 121.4±151.2 | 87.0±37.1 | 0.058 |
| Foot-right | 131.8±176.3 | 84.7±36.1 | 0.025 |

Values are presented as mean ± standard deviation.

Threshold values were taken from the median nerve, on the index finger (hand) and deep and superficial peroneal nerves on the first toe (foot).

DPN, diabetic peripheral neuropathy.