

**Supplementary Table 4.** Univariate analyses of the associations of quadruple therapy discontinuation with various clinical parameters

Variable	Univariate analysis	
	Log-rank statistics	P value
Age, yr	1.007 (0.987–1.027)	0.514
Male sex	0.945 (0.613–1.456)	0.796
BMI, kg/m <sup>2</sup>	0.927 (0.845–1.017)	0.107
Duration of diabetes, yr	0.987 (0.956–1.018)	0.406
HbA1c, %	1.132 (0.979–1.310)	0.095
Alcohol		
Current-drinker	0.644 (0.332–1.248)	0.192
Ex-drinker	1.895 (0.890–4.033)	0.097
Smoking		
Current-smoker	1.188 (0.536–2.634)	0.671
Ex-smoker	1.538 (0.803–2.947)	0.194
Hypertension	0.978 (0.632–1.516)	0.922
Dyslipidemia	1.115 (0.661–1.881)	0.682
Diabetic retinopathy	0.892 (0.534–1.488)	0.661
Diabetic neuropathy	0.758 (0.465–1.234)	0.265
Coronary artery disease	1.396 (0.809–2.408)	0.231
Cerebral artery disease	1.254 (0.627–2.508)	0.522
Peripheral artery disease	1.487 (0.544–4.069)	0.439

Values are presented as hazard ratio (95% confidence interval).  
 BMI, body mass index; HbA1c, glycosylated hemoglobin.