

Supplementary Table 8. Mean glucose levels according to hospital days in patients treated with conventional BBI vs. structured BBI in period 2

	Estimated means, mg/dL		Adjusted <i>P</i> value
	Conventional BBI	Structured BBI	
Fasting			
Day 2	165.9±7.5	174.0±6.0	0.401
Day 3	161.5±6.8	168.2±5.5	0.444
Day 4	147.7±5.9	149.8±4.8	0.782
Day 5	139.5±5.2	137.0±4.3	0.707
Day 6	127.1±4.9	132.1±4.2	0.438
Total	148.2±4.3	152.5±3.5	0.432
Preprandial			
Day 1	324.4±17.4	314.6±14.0	0.640
Day 2	200.0±6.0	192.0±4.9	0.297
Day 3	184.1±5.1	182.4±4.2	0.787
Day 4	173.5±8.5	170.6±7.8	0.624
Day 5	153.4±4.6	159.6±3.8	0.295
Day 6	157.3±4.2	152.0±3.6	0.334
Total	177.7±3.7	177.5±3.0	0.956

Values are presented as mean ± standard error. Analysis of covariance (ANCOVA) analyses were performed, adjusted for age, sex, body mass index, weight, estimated glomerular filtration rate, diabetes duration, glycated hemoglobin, number of concurrent antidiabetic medications, antidiabetic medication use, and number of plasma glucose measurements by point-of-care testing.

BBI, basal-bolus insulin.