| Study or Subgroup | Dip Events | Total | No dip Events | Total | Weight | Odds Ratio MH, Random, 95% Cl | | Ratio om, 95% Cl |
|--|---------------|-------|------------------|-------|--------|----------------------------------|----------------|---------------------|
| eGFR dip 0–10% | | | | | | | | |
| Cherney 2022 | 70 | 1655 | 29 | 752 | 29.9% | 1.101 [0.708; 1.712] | | |
| Oshima 2021 | 35 | 600 | 18 | 294 | 16.9% | 0.950 [0.528; 1.707] | | |
| Total (95% CI) | | 2255 | | 1046 | 46.8% | 1.044 [0.424; 2.572] | | |
| Heterogeneity: $Tau^2 = 0$; $Chi^2 = 0.16$, $df = 1$ (P = 0.69); $l^2 = 0\%$ | | | | | | | | |
| eGFR dip >10% | 6 | | | | | | | |
| Cherney 2022 | 102 | 2141 | 29 | 751 | 32.8% | 1.245 [0.817; 1.898] | | 1 |
| Oshima 2021 | 71 | 956 | 18 | 294 | 20.4% | 1.230 [0.721; 2.100] | | - |
| Total (95% CI) | | 3097 | | 1045 | 53.2% | 1.240 [1.148; 1.338] | | • |
| Heterogeneity: $Tau^2 = 0$; $Chi^2 = 0$, $df = 1$ (P = 0.97); $I^2 = 0\%$ | | | | | | | | |
| Total (95% CI) | | 5352 | | 2091 | 100.0% | 1.144 [0.954; 1.371] | - | - |
| Heterogeneity: Tau ² = 0; Chi ² = 0.64, df = 3 (P = 0.89); $I^2 = 0\%$ | | | | | | | | |
| Test for overall effect: $t_3 = 2.36$ (P = 0.10) | | | | | | | 0.5 | 1 2 |
| Test for subgroup differences: $Chi^2 = 5.83$, df = 1 (P = 0.02) | | | | | | | Favors dipping | Favors no dipping |

Supplementary Fig. 11. Forest plot showing the comparison in hyperkalemia between the dipping and non-dipping groups. MH, Mantel-Haenszel; CI, confidence interval; eGFR, estimated glomerular filtration rate.