Study or Subgroup	Dip Events	Total	No dip Events	Total	Weight	Odds Ratio MH, Random, 95% Cl	Odds Ratio MH, Random, 95% Cl
eGFR dip 0-10)%						
Cherney 2022	585	1655	258	752	28.6%	1.047 [0.873; 1.255]	
Jongs 2022	163	601	69	224	8.3%	0.836 [0.598; 1.170]	
Oshima 2021	188	600	99	294	10.6%	0.899 [0.668; 1.210]	
Total (95% CI)		2856		1270	47.5%	0.973 [0.733; 1.291]	
Heterogeneity:	$Tau^2 = 0;$	Chi ² =	1.68, df	= 2 (P :	= 0.43); l ²	= 0%	
eGFR dip >109	%						
Cherney 2022	738	2141	257	751	30.6%	1.011 [0.849; 1.205]	
Jongs 2022	317	1026	69	224	9.6%	1.004 [0.734; 1.374]	
Oshima 2021	322	956	99	294	12.3%	1.000 [0.759; 1.319]	
Total (95% CI)		4123		1269	52.5%	1.007 [0.993; 1.021]	•
Heterogeneity:	$Tau^2 = 0;$	Chi ² =	0, df = 2	(P = 1.	.00); $I^2 = 0$)%	
Total (95% CI)		6979		2539	100.0%	0.991 [0.918; 1.070]	-
Heterogeneity: Tau ² = 0; Chi ² = 1.81, df = 5 (P = 0.87); $I^2 = 0\%$							
Test for overall effect: $t_5 = -0.31$ (P = 0.77)							0.75 1 1.5
Test for subgroup differences: $Chi^2 = 0.28$, df = 1 (P = 0.60)							Favors dipping Favors no dippin

Supplementary Fig. 7. Forest plot depicting the comparison of serious adverse events between the dipping and non-dipping groups. MH, Mantel-Haenszel; CI, confidence interval; eGFR, estimated glomerular filtration rate.