

SUPPLEMENTARY REFERENCES

1. Park SJ, Choi HR, Choi JH, Kim KW, Hong JP. Reliability and validity of the Korean version of the patient health Questionnaire-9 (PHQ-9). *Anxiety Mood* 2010;6:119-24.
2. Spitzer RL, Kroenke K, Williams JB, Lowe B. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med* 2006;166:1092-7.
3. Pfizer Inc. The Korean version of the GAD-7. Available from: <https://www.phqscreeners.com> (cited 2023 Oct 25).
4. Cohen S, Kamarck T, Mermelstein R. A global measure of perceived stress. *J Health Soc Behav* 1983;24:385-96.
5. Park JO, Seo YS. Validation of the perceived stress scale (PSS) on samples of Korean university students. *Korean J Psychol Gen* 2010;29:611-29.
6. Lee EH. Review of the psychometric evidence of the perceived stress scale. *Asian Nurs Res* 2012;6:121-7.
7. Fisher L, Glasgow RE, Mullan JT, Skaff MM, Polonsky WH. Development of a brief diabetes distress screening instrument. *Ann Fam Med* 2008;6:246-52.
8. Choi S, Kim SH. Influences of patient activation on diabetes self-care activities and diabetes-specific distress. *Korean J Adult Nurs* 2020;32:10-20.
9. Nam HS, Kim KY, Kwon SS, Koh GW, Kind P. Report on the estimation of quality weights for the European Quality of Life-5 Dimensions (EQ-5D). Cheongju: Korea Disease Control and Prevention Agency; 2007.
10. Fitzgerald JT, Davis WK, Connell CM, Hess GE, Funnell MM, Hiss RG. Development and validation of the diabetes care profile. *Eval Health Prof* 1996;19:208-30.
11. Rhee CW. Relationship between diabetes-related social support and depression in people with diabetes. *Korean J Soc Welf Stud* 2007;33:273-93.
12. Seo K, Song M, Choi S, Kim SA, Chang SJ. Development of a scale to measure diabetes self-management behaviors among older Koreans with type 2 diabetes, based on the seven domains identified by the American Association of Diabetes Educators. *Jpn J Nurs Sci* 2017;14:161-70.