

Supplementary Table 2. Distribution of adults eligible for statin therapy in each year's Korean National Health and Nutrition Examination Surveys from 1998 to 2012

Variable	I	II	III	IV			V		
	1998	2001	2005	2007	2008	2009	2010	2011	2012
ASCVD	1.4 (0.1)	1.6 (0.2)	3.5 (0.3)	3.9 (0.5)	3.4 (0.3)	2.6 (0.2)	3.0 (0.3)	3.5 (0.3)	3.4 (0.3)
LDL-C \geq 190 mg/dL	1.7 (0.2)	1.2 (0.2)	1.1 (0.2)	1.3 (0.2)	1.6 (0.2)	1.4 (0.2)	1.5 (0.2)	1.6 (0.2)	1.7 (0.2)
Diabetes mellitus	4.8 (0.3)	5.6 (0.3)	4.9 (0.3)	5.3 (0.5)	5.7 (0.3)	5.5 (0.3)	5.6 (0.4)	6.7 (0.4)	6.1 (0.4)
10-Year risk \geq 7.5%	9.1 (0.4)	10.6 (0.5)	11.3 (0.5)	9.2 (0.6)	9.5 (0.5)	10.9 (0.5)	11.3 (0.6)	10.5 (0.6)	11.1 (0.5)
Adults eligible for statin therapy	17.0 (0.7)	19.0 (0.7)	20.8 (0.7)	19.6 (1.1)	20.2 (0.8)	20.4 (0.8)	21.4 (0.9)	22.2 (0.9)	22.3 (0.8)
Adults taking lipid-lowering therapy	NA	NA	1.0 (0.1)	1.8 (0.3)	2.4 (0.2)	3.0 (0.2)	4.2 (0.3)	4.3 (0.3)	5.3 (0.4)

Values are presented as percentage (standard error).

ASCVD, atherosclerotic cardiovascular disease; LDL-C, low density lipoprotein cholesterol; NA, not assessed.