

Supplementary Table 3. Distribution of adults eligible for statin therapy according to sex and age in each year's Korean National Health and Nutrition Examination Surveys from 1998 to 2012

Age, yr	Men										Women									
	I		II		III		IV		V		I		II		III		IV		V	
	1998	2001	2005	2007	2008	2009	2010	2011	2012	1998	2001	2005	2007	2008	2009	2010	2011	2012		
20-29	0.7 (0.3)	0.9 (0.5)	0.0	0.5 (0.5)	1.9 (0.8)	0.0	1.1 (0.5)	0.6 (0.6)	0.2 (0.2)	1.5 (0.5)	0.5 (0.3)	0.1 (0.1)	1.5 (0.9)	0.2 (0.2)	0.8 (0.4)	0.5 (0.3)	0.6 (0.6)	0.4 (0.3)		
30-39	1.9 (0.5)	1.1 (0.5)	0.8 (0.4)	1.3 (0.8)	1.6 (0.6)	1.2 (0.5)	2.7 (0.9)	1.7 (0.7)	2.8 (1.0)	0.5 (0.2)	1.1 (0.5)	0.8 (0.4)	0.5 (0.3)	1.0 (0.5)	1.4 (0.5)	1.1 (0.6)	1.3 (0.6)	1.7 (0.7)		
40-49	19.4 (1.8)	23.9 (2.0)	21.4 (2.0)	14.7 (2.7)	18.3 (1.9)	17.5 (1.8)	17.3 (2.1)	19.0 (2.3)	14.8 (2.0)	6.6 (1.0)	5.6 (0.9)	7.2 (1.3)	4.5 (1.4)	6.6 (1.0)	7.4 (1.0)	6.4 (1.5)	8.9 (1.4)	7.7 (1.3)		
50-59	56.5 (2.4)	56.2 (3.2)	59.3 (2.7)	49.1 (4.0)	45.4 (3.0)	48.1 (2.3)	49.2 (2.7)	49.5 (3.0)	51.2 (2.8)	19.9 (1.8)	18.9 (2.3)	20.2 (2.2)	19.9 (2.8)	17.3 (1.7)	13.8 (1.4)	16.7 (1.7)	15.4 (1.7)	17.3 (1.7)		
60-69	79.7 (2.2)	92.9 (1.5)	91.1 (1.6)	88.0 (2.6)	83.3 (2.5)	87.9 (1.8)	88.5 (1.8)	88.4 (1.8)	90.3 (1.6)	53.8 (2.6)	56.5 (3.0)	53.3 (2.7)	50.1 (4.1)	49.8 (2.3)	49.4 (2.5)	49.2 (2.7)	52.2 (2.5)	47.7 (2.6)		
70-79 ^a	60.1 (4.3)	79.2 (3.9)	74.3 (4.5)	70.2 (4.6)	71.0 (3.1)	71.0 (2.8)	74.2 (3.6)	74.5 (2.7)	68.9 (3.6)	65.9 (3.3)	66.9 (3.7)	73.0 (3.3)	69.7 (4.6)	67.7 (3.0)	67.1 (2.5)	68.3 (3.2)	63.5 (3.0)	63.3 (3.3)		
70-75	82.3 (4.5)	95.3 (2.5)	94.6 (2.5)	94.8 (1.8)	90.4 (2.4)	90.2 (1.9)	92.3 (2.3)	95.4 (1.4)	93.9 (2.6)	88.9 (2.4)	95.6 (1.8)	96.1 (1.6)	93.2 (2.6)	94.0 (1.7)	94.2 (1.4)	95.2 (1.5)	93.9 (1.5)	92.3 (1.8)		
76-79	3.4 (2.3)	19.2 (7.8)	2.7 (2.7)	8.3 (5.0)	21.4 (5.5)	15.3 (4.5)	12.9 (4.8)	23.1 (5.0)	19.4 (4.6)	9.5 (4.1)	5.5 (3.0)	14.4 (3.9)	26.6 (7.2)	17.3 (4.4)	14.8 (3.3)	16.1 (4.8)	11.5 (2.9)	10.1 (2.6)		
≥80	13.6 (8.8)	6.1 (4.8)	5.7 (4.0)	19.2 (9.9)	12.6 (5.3)	19.8 (5.2)	18.3 (7.3)	28.3 (9.4)	9.7 (6.9)	10.8 (4.6)	3.8 (2.2)	20.4 (6.2)	12.9 (4.5)	16.5 (4.9)	5.0 (2.9)	7.3 (3.7)	26.3 (5.2)	15.4 (4.9)		
Percentage of adults aged ≥20 years eligible for statin therapy, %	20.7 (0.9)	24.3 (1.1)	25.8 (1.1)	24.0 (1.3)	25.0 (1.1)	25.6 (1.0)	27.2 (1.2)	28.0 (1.4)	27.8 (1.2)	13.5 (0.7)	13.9 (0.8)	16.1 (0.8)	15.4 (1.1)	15.6 (0.7)	15.4 (0.8)	15.8 (0.9)	16.6 (0.8)	16.8 (0.9)		

Values are presented as percentage (standard error).

^aThe 70- to 79-year-old group was divided into two groups, 70 to 75 and 76 to 79 years, to assess the percentages of people classified into the statin benefit group before and after the age of 75 years because the third and fourth criteria of the 2013 American College of Cardiology/American Heart Association guideline limits age to 40 to 75 years.