

Supplementary Table 1. Weighted percent of the all-cause and cause-specific mortality rate according to the glycemic control status of diabetes ($n=2,391$)

Variable	Overall		Glycemic control status				P value
	No. (%)	SE	Yes		No		
			No. (%)	SE	No. (%)	SE	
Deaths							
All-cause	478 (18.27)	0.99	422 (18.27)	1.09	56 (18.23)	2.73	0.99
CVD	100 (3.87)	0.51	87 (3.96)	0.64	13 (3.37)	1.21	0.71
Cancer	81 (3.01)	0.46	65 (2.55)	0.48	16 (5.58)	1.50	0.02
CLRD	25 (1.14)	0.24	24 (1.27)	0.28	1 (0.42)	0.42	0.25
Accidents	8 (0.21)	0.10	8 (0.25)	0.12	0	0.00	NA
CeVD	28 (1.33)	0.27	23 (1.32)	0.30	5 (1.38)	0.71	0.94
AD	5 (0.19)	0.10	5 (0.22)	0.11	0	0.00	NA
Flu & pneumonia	18 (0.58)	0.22	15 (0.47)	0.15	3 (1.23)	1.02	0.22
Kidney disease	19 (0.51)	0.16	17 (0.50)	0.16	2 (0.57)	0.42	0.85

Glycemic control status of diabetes was inferred by medication (insulin or diabetic pills) usage status.

SE, standard error; CVD, cardiovascular disease; CLRD, chronic lower respiratory disease; NA, not available; CeVD, cerebrovascular disease; AD, Alzheimer's disease; Flu, influenza.